species tend to be restricted to lowland forest, montane forest or the subalpine zone. Powelliphanta

Powelliphantæspecific primer (POWCO1r) that replaces H7005 (Hafner et al. 1994) used in previous studies. Together with LCO1490 (Folmer et al. 1994), POWCO1r (5′ GCA ATT ACT ATC GTT GCA GCA G 3′) targets a fragment of about 700 bp at the 5′ end of COI. PCR reactions were performed in 10 µL volumes with Red Hot® Taq polymerase using thermocycling conditions 94°C for

calculated and log transformed as described above. Loci were examined separately and as a single group.

We searched for evidence of population structuring among our multi-locus genotype data using Bayesian assignment in STRUCTURE v2.3.2 (Pritchard et al. 2000). This approach infers genotype similarity and assigns individuals to clusters (populations), as well as determining isolation-by-barrier. We used a Bayesian Markov Chain Monte Carlo (MCMC) simulation chain of 100,000 steps following a burnin of 10,000 steps. The number of clusters (K) tested was set at between 1 and 15 in separate runs with 11 iterations of each. This analysis was performed using the no-admixture model, appropriate for studying fully discrete populations and suitable for detecting subtle structure. Analysis was repeated with both correlated and independent allele frequency models and then repeated using the admixture model. We also examined the e ect of including sampling location information as recommended for analyses with relatively small sample sizes (Hubisz et al. 2009). Including sampling location categories or phenotype-based groups allows the models implemented by STRUCTURE v2.3.2 to modify the prior distribution for each individual's population assignment. Results were processed in STRUCTURE HARVESTER v0.6.93 (Earl and vonHoldt 2011) to assess and visualise likelihood values across multiple values of K and hundreds of iterations to determine the number of genetic clusters that best fitted the genotypic data. This used the DeltaK statistic that considers the rate of change in log probability between successive values of K. In addition, STRUCTURE HARVESTER v0.6.93 was used to reformat data for optimal K for analysis in CLUMPP v1.1.2 (Jakobsson and Rosenberg 2007). This cluster matching and permutation program deals with label switching and multimodality in analysis of population structure. DISTRUCT v1.1 (Rosenberg 2003) was used to illustrate results.

To determine which populations lent significance to the genetic structure of the group we used OBSTRUCT (Gayevskiy et al. 2014). CSV files were manually created from STRUCTURE output files for hypothesised numbers of populations (U(i)13.2(n)763.3(fo)43



Figure 2. Population genetic structure among land snail populations in the New Zealand Powelliphanta Kawatiri complex. (A) Population sampling for mtDNA haplotypes; (B) median joining network of mtDNA COI (622bp) haplotypes with colours indicating geographic source shown on map; (C) population sample clusters indicated by Bayesian genotype assignment of ve microsatellite loci shown in graphs; (D) no-admixture model, K = 2; (E) admixture model, K = 2; (F) non-admixture model, K = 6; (G) admixture model, K = 6. Map C shows population samples sharing genotypic clusters (dashed lines). Colours represent alternative genotype assignments, at each K.

departure from the expectations of the Hardy-Weinberg equilibrium for four of the five loci within the population sample at Stockton, probably caused by the presence of null alleles. Pairwise linkage disequilibrium tested using the genotypes of the 31 Stockton snails indicated that the alleles at di erent loci randomly associated. The number of alleles per locus ranged from 4 to 20. Per locus, allelic richness from all genotyped population samples (n = 15) reached eight (Supplementary Table S3). Average gene diversity (Table 2) was highest in the sample from Brunner Mid (0.733). Braeburn and Matakitaki samples had no allelic variation, but each consisted of only two snails. Allele sharing among population samples was common but half of the population pairwise F_{ST} estimates were significantly greater than zero (46/91; Table S4). Each population was significantly di erentiated from at least three other populations, with the exception of Brunner North (n = 2; Supplementary Table S3). Stockton di ered significantly from all other population samples (pairwise F_{ST} 0.247–0.504). Even Denniston South and Denniston North di ered significantly although the collection sites of these samples were only 5.4 km apart. Microsatellite di erentiation followed a model of isolation by distance (Mantel test: r = 0.3207; P = .0190).

Population structure

Bayesian assignment of microsatellite genotypes using STRUCTURE v2.3.2 yielded similar results using correlated and independent allele frequency models. The maximum value of Delta K determined using the independent allele frequency model, which is reported as less likely to overestimate K, was for K = 2 with the next optimal value for K = 6. Visual examination of output from DISTRUCT (Figure 2) shows that K = 6 cap-

are two major groupings seen within the genetic diversity of the Kawatiri complex lineage, one comprises the populations found on the western edge of their range (Paparoa, Mount William and Brunner ranges), and the other includes the more inland (eastern) populations with six shallower clusters. We advocate management that retains the evolutionary potential of these clusters